

## INSTRUCTIONS FOR USE

CONTENTS: Emtrix, 10 ml

### INGREDIENTS

Propylene Glycol, Urea (20%), Glycerol, Lactic Acid (9%), Water, Sodium Hydroxide, Disodium EDTA

### DESCRIPTION

Emtrix is a clear, colourless liquid supplied in a plastic tube with a silicone drop tip to facilitate application to the nail. Emtrix is free from preservatives and fragrances.

### INDICATION

For treatment of nails affected by fungal infection (onychomycosis) or psoriasis.

Emtrix improves the appearance of discoloured and deformed nails. Emtrix has a physical keratolytic effect and degrades the cells of fungi that typically cause fungal nail infection.

Emtrix increases hydration, gently smoothes the outer layers of the nail plate and, by changing the nail's microenvironment and improving nail surface integrity, keeps nail in good condition. First signs of improvement are normally seen after one to two weeks of treatment. In a clinical study 76% of the users saw improvement in nail appearance in just 1 week, and 92% after 8 weeks\*.

### USE

Apply Emtrix once daily to affected nails, preferably in the evening just before bed time. Hold the point of the tube onto the nail. Press lightly on the tube and spread out the solution with the point. Cover each nail with a thin layer of the solution. It is important that the solution is also applied under the free edge of any affected nails. After application, the nail should be allowed to dry for a few minutes.

If you use nail polish during treatment with Emtrix, remove any old layers of nail polish thoroughly in the evening. Apply Emtrix to affected nails before going to bed, as described above. The next day, nail polish can be applied to the treated nails. Repeat this procedure every day when using nail polish during treatment. Duration of treatment varies according to the initial severity of the condition, but in many cases 3-6 months treatment may be required. For optimal results, continue the treatment until a normal nail has grown out.

Do not apply Emtrix to the nail bed if the affected nail or parts of the affected nail has detached from the underlying nail bed. Emtrix acts only locally and can therefore be used during pregnancy and lactation. There is limited experience in children. Therefore, consult your doctor before treatment in children.

### WARNINGS AND PRECAUTIONS

For external use only. Avoid contact with eyes or mucous membranes. Do not use if you are allergic to any of the ingredients.

### POSSIBLE SIDE EFFECTS

Transient minor irritation of the skin immediately adjacent to the nail and discolouration of the nail can be expected to occur in some patients. In addition, loosening/separation of the nail from the underlying nail bed (onycholysis) may occur, including detachment of the affected part of the nail.

Any serious incident that has occurred in relation to the device should be reported to the manufacturer and the competent authority of the Member State in which you live in.



### SHELF LIFE AND STORAGE

Use by the date indicated on the tube and outer carton. Store between 2 °C and 25 °C. Protect from sunlight and freezing. Keep out of reach of children.

Advantice Health, LLC.  
685 Route 202/206  
Bridgewater  
New Jersey  
08807  
USA

DISTRIBUTOR  
Advantice Health AB  
c/o Covendum  
Biblioteksgatan 29  
SE-114 35 Stockholm  
Sweden

EMTRIX® is a registered trademark of Advantice Health AB.



### MedEnvoy, Global B.V.

Prinses Margrietplantsoen 33 - Suite 123  
2595 AM The Hague  
The Netherlands



Date of issue: 2024-02, ver 04